



Role of Confounding Factors in the Evaluation of Vitamin D Deficiency

Masoud Nouri-Vaskeh¹ , Elaheh Ouladsahebmadarek^{2*} 

Dear Editor,

Vitamin D has been traditionally used as a mineral regulator in preventing musculoskeletal disorders. Nowadays, the beneficial effects of vitamin D on the skeletal system and several body organs have been identified (1, 2). Recent studies have asserted that vitamin D contributes to the processes of anti-inflammatory and pro-inflammatory cytokines, as well as suppression of angiogenesis (3). The correlation between vitamin D level and several diseases has been investigated in numerous studies, however only a few have exactly considered the role of peripheral confounding factors on the levels of vitamin D. In the current letter, we aimed to define the important associated factors that should be taken into account in the determination of vitamin D deficiency in all patients and diseases in all regions.

Sunlight plays a critical role in vitamin D metabolism. It differs in various regions of the earth's surface, thereby making a difference in the amount of sunlight exposure. In addition, the sunlight exposure is important in the synthesis of endogenous vitamin D. However, inadequate exposure due to women's extensive coverage for religious or cultural reasons, for example, can affect vitamin D status. Furthermore, in some regions with hot climate, women have to stay at home during the day, and as a result, they will be divested of enough sunlight exposure (4). Moreover, the effects of pregnancy, ethnicity, skin color, and genetic background on vitamin D status are well-clarified (5). Recent studies have focused on aging and high prevalence of vitamin D deficiency among centenarians (6). Additionally, the effect of sunlight in various seasons, especially in winter, must be taken into consideration (7). In addition to parathyroid disorders, other comorbidities including chronic kidney or liver diseases play a crucial role in the serum levels of vitamin D (8, 9). The dietary habits such as eating a great deal of marine food can increase the vitamin D level; in contrast, malnutrition reduces its concentration (10, 11).

The 25(OH)D assay is a routine method for the diagnosis

of vitamin D deficiency, and the stability of 25(OH)D is more than 1,25(OH)₂D owing to the longer half-life and higher concentration of 25(OH)D about 1000 fold (12).

The serum concentration of 25(OH)D is affected by thyroid disorders, as well as some drugs such as antiepileptics and cholestyramine. The serum levels of 1,25(OH)₂D can also be influenced by phosphate and calcium, as well as sedentary lifestyle. Moreover, the parathyroid hormone and its peptides, prostaglandins, estradiol, and prolactin have an influence on the metabolism of 1,25(OH)₂D. The effect of drugs including heparin sodium, corticosteroids, statins, antihypertensives, anticonvulsants, anti-retroviral drugs, and bisphosphonates on 1,25(OH)₂D is indisputable (13). Higher body mass index is also an important confounding factor associated with incomplete response to vitamin D supplement therapy and elevation of 25(OH)D is less than expected (10). In addition, division of vitamin D deficiency into the groups of insufficiency and deficiency will make it easy to interpret the results.

Overall, considering the confounding factors with the ability of changing the levels of vitamin D is of great importance.

Conflict of Interests

None declared.

Ethical Issues

Not applicable.

Financial Support

No funding was received for this study.

References

1. Hossein-Nezhad A, Holick MF. Vitamin D for health: a global perspective. *Mayo Clin Proc.* 2013;88(7):720-755. doi:10.1016/j.mayocp.2013.05.011
2. Derakhshan M, Derakhshan M, Hedayat P, Shiasi M, Sadeghi E. Vitamin D Deficiency May Be a Modifiable Risk



- Factor in Women With Endometriosis. *Crescent Journal of Medical and Biological Sciences*. 2018;5(4):292-296.
3. Calton EK, Keane KN, Soares MJ. The potential regulatory role of vitamin D in the bioenergetics of inflammation. *Curr Opin Clin Nutr Metab Care*. 2015;18(4):367-373. doi:10.1097/mco.000000000000186
 4. Iftikhar R, Kamran SM, Qadir A, Haider E, Bin Usman H. Vitamin D deficiency in patients with tuberculosis. *J Coll Physicians Surg Pak*. 2013;23(10):780-783. doi:11.2013/jcsp.780783
 5. Herrmann M, Farrell CL, Pusceddu I, Fabregat-Cabello N, Cavalier E. Assessment of vitamin D status - a changing landscape. *Clin Chem Lab Med*. 2017;55(1):3-26. doi:10.1515/cclm-2016-0264
 6. Yao Y, Fu S, Li N, et al. Sex, Residence and Fish Intake Predict Vitamin D Status in Chinese Centenarians. *J Nutr Health Aging*. 2019;23(2):165-171. doi:10.1007/s12603-018-1126-1
 7. Levis S, Gomez A, Jimenez C, et al. Vitamin d deficiency and seasonal variation in an adult South Florida population. *J Clin Endocrinol Metab*. 2005;90(3):1557-1562. doi:10.1210/jc.2004-0746
 8. Nair S. Vitamin d deficiency and liver disease. *Gastroenterol Hepatol (N Y)*. 2010;6(8):491-493.
 9. Williams S, Malatesta K, Norris K. Vitamin D and chronic kidney disease. *Ethn Dis*. 2009;19(4 Suppl 5):S5-8-11.
 10. Mazahery H, von Hurst PR. Factors Affecting 25-Hydroxyvitamin D Concentration in Response to Vitamin D Supplementation. *Nutrients*. 2015;7(7):5111-5142. doi:10.3390/nu7075111
 11. Hosseinzadeh F, Jangi Oskouei N, Ghavamzadeh S. The Effect of Vitamin D Supplementation on Inflammation and Markers of Vascular Function in Heart Failure Patients. *Crescent Journal of Medical and Biological Sciences*. 2019;6(3):285-292.
 12. Lips P. Relative value of 25(OH)D and 1,25(OH)2D measurements. *J Bone Miner Res*. 2007;22(11):1668-1671. doi:10.1359/jbmr.070716
 13. Gröber U, Kisters K. Influence of drugs on vitamin D and calcium metabolism. *Dermatoendocrinol*. 2012;4(2):158-166. doi:10.4161/derm.20731

Copyright © 2019 The Author(s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.