



# The Relationship Between Scapular Dyskinesia and Generalized Joint Hypermobility in Young Women

Afsun Nodehi Moghadam<sup>1\*</sup>, Maryam Moghadam Salimi<sup>2</sup>, Enayatollah Bakhshi<sup>3</sup>

## Abstract

**Objectives:** Individuals with generalized joint hypermobility (GJH) have motions beyond the normal range in the shoulder, and less shoulder stability. Scapular dyskinesia is likely to be a consequence of weakness in scapular stabilizer muscle. This study intended to identify the relationship between GJH and scapular dyskinesia in young women.

**Materials and Methods:** One hundred women (47 hypermobile and 53 non-hypermobile) participated in this case control study. The Beighton score was used to diagnose GJH. Visual scapular dyskinesia test (proposed by Uhl et al) was used for the evaluation of scapular dyskinesia. Scapular winging and/or dysrhythmia is observed during a set of bilateral, active, shoulder flexion, abduction and scaption (40° anterior to the frontal plane) for 5 times while having the thumbs pointed up. These movements were performed, bearing the weight on their hands. Generalized estimating equations were used in order to compare the scapular dyskinesia prevalence during shoulder flexion, abduction and scaption between the females with and without GJH.

**Results:** The prevalence of scapular dyskinesia in the females with GJH was twice more than that in the females with no hypermobility [OR=2.18(95% CI: 1.18-4.03)]. It was also found that there were not any significant differences in the prevalence of scapular dyskinesia between shoulder elevation planes (flexion, abduction and scaption) in the females with and without GJH ( $P>0.05$ ).

**Conclusions:** Higher prevalence of scapular dyskinesia in the females with GJH may place them at the risk of future shoulder pain and pathology, which should be considered in the evaluation and management of hypermobile individuals.

**Keywords:** Joint hypermobility, Joint laxity, Scapula, Shoulder injuries

## Introduction

Generalized joint hypermobility (GJH) is a term used when most of the synovial joints in the body have an increased range of motion due to excessive connective tissue extensibility and/ or capsular or ligamentous laxity (1,2). GJH is a feature present in various rheumatologic diseases, including Marfan syndrome and Ehlers-Danlos syndrome, but it is known as a benign joint hypermobility syndrome that can also be present in the absence of rheumatologic diseases (3,4). GJH is most often seen in the females and among African and Asian descents, and its prevalence rate is reported from 5% to 43 % (5,6). Several studies have shown that the individuals with GJH have widespread joint pain and musculoskeletal and non-musculoskeletal complaints (1,,7-9). Approximately 3.3% of females suffering from GJH have tendency to develop pain in the legs and shoulder joint (10) . The result of Trudelle-Jackson and colleagues' study showed that hypermobility appears to place women at increased risk of musculoskeletal injury while participating in physical

activity (11).

In the individuals with GJH, lower joint stability along with muscle weakness can be a major risk factor in the development of upper extremity disorders (12) .

Arm elevation requires coordinated motions of glenohumeral and scapulothoracic joints (13). Shoulder girdle stability is provided by static stabilizers (such as capsule, ligaments), dynamic stabilizers (including rotator cuff and scapular stabilizers), as well as proprioceptive information emerging from mechanoreceptors in muscles, tendons, joint-capsule ligaments, and skin, which are centrally integrated (13). Optimal scapular position and motion depend on normal function of the scapular stabilizer muscles such as trapezius, rhomboids, and serratus anterior muscles (13).

Scapular dyskinesia is a change from normal scapular position and motion and is characterized by prominence of scapular medial border and/or inferior angle, early scapula elevation or shrugging and reduced upward rotation of the scapula during arm elevation (14). Scapular

Received 5 June 2017, Accepted 19 December 2017, Available online 10 January 2018

<sup>1</sup>Department of Physiotherapy, Rofeideh Rehabilitation Hospital, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran. <sup>2</sup>Department of Physiotherapy, Tabriz University of Medical Sciences, Tabriz, Iran. <sup>3</sup>Department of Statistics, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran.

\*Corresponding Author: Afsun Nodehi Moghadam, PhD PT; Tel: (98)-21-22180084, Email: [afsoonnodehi@gmail.com](mailto:afsoonnodehi@gmail.com)



dyskinesia is observed among the individuals suffering from various shoulder disorders such as glenohumeral instability, rotator cuff disorders, and labrum tears (15-19). Since scapula has limited bony attachment, its mobility and stability depends on muscle activation especially serratus anterior and upper and lower trapezius muscles (13). Alterations of periscapular muscle activation such as reduced strength of lower trapezius and serratus anterior muscles are common among the individuals with scapular dyskinesia (15). It has also been found that women with GJH show physical fitness at lower levels, poor joint proprioception, and decreased muscle strength (20-22). Thus, since scapular dyskinesia is a consequence of weakness in scapular stabilizer muscle, it is possible that women with GJH have higher prevalence of scapular dyskinesia (12,15). Therefore, this study tried to identify the relationship between GJH and scapular dyskinesia among young women.

## Materials and Methods

### Subjects

One hundred women (47 hypermobile and 53 nonhypermobile) participated in this case control study. The Beighton score was applied to diagnose GJH with a scoring range from 0 to 9. Beighton scoring system consists of four bilateral tests including (a) passive hyperextension of little finger beyond 90°, (b) passive reaching of the thumbs to the flexor aspects of the forearms, (c) elbows hyperextension beyond 10°, (d) knees hyperextension more than 10°, and one unilateral test: (e) trunk flexion so that the palms can easily touch the floor. A cut point of 4-9 or higher can meet the Beighton score for GJH (23). In this study, participants with score of 5 or higher were considered as hypermobile subjects. The participants in non-hypermobile group (with a Beighton score lower than 5) were matched in age, height, and weight with GJH group. The subjects with a history of shoulder pain, dislocation, fracture, or surgery as well as neurologic or rheumatologic diseases and scoliosis or excessive kyphosis were excluded (17). All subjects showed their agreement by signing the consent form which was approved by the Human Subjects Committee of University of Social Welfare and Rehabilitation Sciences.

### Scapular Dyskinesia Test

In this study, visual method (proposed by Uhl et al) was used for the evaluation of scapular dyskinesia (24). The participants stood with their arms resting next to their bodies, while the examiner stood behind them at 1.5-meter distance. Each participant performed a set of bilateral, active, shoulder flexion, abduction and scaption (40° anterior to the frontal plane) for 5 times while having the thumbs pointed up. These movements were performed, bearing the weight on their hands. The participants were required to raise their arms as far as possible over a count of 3 seconds and then descend them over a count

of 3 seconds. The movements were performed while holding dumbbells on the basis of their body weight as 1 kg for less than 60 kg and 2 kg for 60 kg or more (25). A pilot study was performed so as to choose the weights. Presence or absence of scapular dyskinesia (winging and/or dysrhythmia) was observed during bilateral weighted shoulder elevation (24,25).

### Statistical Analysis

The statistical analysis was conducted using SPSS statistical software version 19.0. Based on the Beighton score (A cut point of 5-9 or higher), the participants were grouped as hypermobile or non-hypermobile. Descriptive statistics was expressed as mean and standard deviation (SD) for participants' age, weight, height and Beighton score, or as percentages for the scapular dyskinesia prevalence. Independent *t* test was used in order to compare the participants' characteristics in the 2 groups. Moreover, because of repeated measurements and correlated observations, generalized estimating equations were used to compare the scapular dyskinesia prevalence during shoulder flexion, abduction and scaption between the females with and without GJH. For all statistical analyses, a significance level of 0.05 was used.

### Results

There were no statistically significant differences regarding the participants' age, weight, and height between the two groups with and without hypermobility ( $P > 0.05$ ) (Table 1). The mean (SD) of Beighton score in the females with GJH and control subjects was  $5.87 \pm 0.95$  and  $2.19 \pm 1.37$ , respectively. The prevalence of scapular dyskinesia during shoulder flexion, abduction and scaption in the females with and without GJH are shown in Table 2. During dominant side flexion, scaption and abduction, scapular dyskinesia was seen in 68.1%, 78.7%, and 68.1% of the participants in the hypermobile group, while its prevalence in the control group was 31.9%, 21.3%, and 31.9%, respectively. Moreover, the prevalence of scapular dyskinesia during non-dominant side flexion, scaption and abduction in the females with GJH was more than that in the females with no hypermobility (Table 2).

Furthermore, the results of generalized estimating equations showed that the prevalence of scapular dyskinesia in the females with GJH was twice more than

**Table 1.** Demographic Characteristics of Females With and Without Generalized Joint Hypermobility

| Variables      | Mean $\pm$ SD     |                   | P Value |
|----------------|-------------------|-------------------|---------|
|                | Hypermobile       | Non-hypermobile   |         |
| Age (y)        | 22.11 $\pm$ 2.52  | 22.75 $\pm$ 2.61  | 0.21    |
| Weight (kg)    | 56.27 $\pm$ 6.33  | 56.42 $\pm$ 6.15  | 0.90    |
| Height (cm)    | 161.91 $\pm$ 5.70 | 162.09 $\pm$ 5.62 | 0.87    |
| Beighton score | 5.87 $\pm$ 0.95   | 2.19 $\pm$ 1.37   | 0.00    |

**Table 2.** The Prevalence of Scapular Dyskinesia During Shoulder Flexion, Abduction and Scaption in Dominant and Non-dominant Sides in Females With and Without Generalized Joint Hypermobility

| Scapular Dyskinesis No. (%) |     | Dominant Side |                 | Non-dominant Side |                 |
|-----------------------------|-----|---------------|-----------------|-------------------|-----------------|
|                             |     | Hypermobile   | Non-hypermobile | Hypermobile       | Non-hypermobile |
| Flexion                     | Yes | 32(68.1)      | 28 (52.8)       | 35 (74.5)         | 33 (62.3)       |
|                             | No  | 15(31.9)      | 25 (47.2)       | 12 (25.5)         | 20 (37.7)       |
| Scaption                    | Yes | 37(78.7)      | 25 (47.2)       | 38 (80.9)         | 33 (62.3)       |
|                             | No  | 10(21.3)      | 28 (52.8)       | 9 (19.1)          | 20 (37.7)       |
| Abduction                   | Yes | 32(68.1)      | 29 (54.7)       | 34 (72.3)         | 31 (58.5)       |
|                             | No  | 15(31.9)      | 24 (45.3)       | 13 (27.7)         | 22 (41.5)       |

that in the females with no hypermobility (OR [odds ratio] =2.18 [95% CI: 1.18-4.03]). However, it was found that there were not any significant differences in the prevalence of scapular dyskinesia between shoulder elevation planes (flexion, abduction and scaption) in the females with and without GJH ( $P > 0.05$ ).

### Discussion

As the result of this study showed, the prevalence of scapular dyskinesia in the females with GJH was twice more than that in the females with no hypermobility. The primary characteristic of the females with GJH is the excessive range of joint motions (1). Hypermobile joint is likely to be less stable, to be subject to subluxation or dislocation, and more susceptible to musculoskeletal injuries (26). It has already been proven that GJH is a significant risk factor for traumatic anterior shoulder dislocation and acute and chronic shoulder injuries (7, 27). The result of a longitudinal cohort study in children suggests an association between GJH in childhood and resulting adolescent joint pain, so that they had three times increased risk of inflicting pain at the age of 14 (9). Moreover, the result of Trudelle-Jackson and colleagues' study showed that the women with multiple joints hypermobility who participate at any level of physical activity are more in danger of musculoskeletal injury (11).

The results of some studies have shown that the individuals with GJH have decreased physical fitness and muscle weakness (20,22). The results of Scheper et al (28) and Jindal and colleagues' (29) studies showed decreased grip, shoulder abductor and elbow extensor strengths in the individuals with GJH. On the other hand, Jeremiah and Alexander (30) suggested that the increased laxity along with GJH can lead to poor neuromuscular control (proprioception). The results of some previous studies indicate that the individuals with GJH demonstrate reduced proprioception than those without joint hypermobility (21,31). It seems that since the females with GJH have more range of shoulder motion and reduced joint stability due to lax static stabilizers, they may rely more on muscle force and neuromuscular mechanisms (proprioception) (30). Thus, according to the results of previous studies which have shown the individuals with

GJH have decreased physical fitness and muscle weakness, higher prevalence of scapular dyskinesia in the females with GJH looks reasonable (20,22).

The prevalence of scapular dyskinesia has been observed in 68%-100% of individuals suffering from shoulder disorders, such as rotator cuff tears, glenohumeral instability, and labral tears (17,32). The findings of this study regarding the association of scapular dyskinesia with GJH are in line with the results of shoulder instability studies (7,27). In agreement with our result, it was shown that scapular dyskinesia was often seen in unstable glenohumeral joint such as multidirectional instability and recurrent types of shoulder instability (14,33,34). It has been suggested that the lax capsular tissue and altered biomechanics and muscle activation in the individuals with multidirectional instability may lead to scapular dysrhythmia, excessive protraction of scapula and so-called scapular dyskinesia (14). Nodehi Moghadam and Salimee (35) showed that compared to non-hypermobile subjects, females with GJH exhibited a more protracted and downward-rotated scapula on their dominant side with the arm at rest beside the body. Moreover, in agreement with our result, Alibazi et al (36) investigated the impact of shoulder muscle fatigue on shoulder kinematic in the females with GJH and found that before fatigue, 55% of GJH women had subtle or obvious scapular dyskinesia while, in control group it was 31.3%. Another mechanism that may justify higher prevalence of scapular dyskinesia in the females with GJH is abnormal neuromuscular control (21). The presence of connective tissue laxity and poor proprioception in the females with GJH may result in compensatory strategies such as co-contraction or excessive activation of specific shoulder girdle stabilizer muscles, which could lead to muscles fatigue and weakness, altered scapulohumeral rhythm and scapular dyskinesia (14,28). The evidence shows that upper limb muscle fatigue can result in scapular dyskinesia (37,38). It has been suggested by some researchers that fatigue-induced scapular dyskinesia may be an etiological factor in the reduction of subacromial space and impingement syndrome (38). While, some others have shown that alteration in scapular orientation has minimal effect on subacromial space width; thus, can be considered as a

compensatory motion to offset superior humeral head translation, subsequent to rotator cuff fatigue (37,39).

This study has some limitations. First, in our study only a homogeneous group of healthy and non-symptomatic women was assessed. Considering such a limitation, applying these findings to the symptomatic and male population should be proceeded with caution; so more studies in these areas are necessary. Second, the probable weakness of scapula muscles in the individuals with scapular dyskinesis as assessed using strength tests or electromyography data was not included in this study.

### Conclusions

According to the results of this study, the prevalence of scapular dyskinesis in the females with GJH was twice more than that in the females with no hypermobility (OR=2.18 [95% CI: 1.18-4.03]). It was also found that there were not any significant differences in the prevalence of scapular dyskinesis between shoulder elevation planes (flexion, abduction and scaption) in the females with and without GJH.

### Conflict of Interests

None declared.

### Ethical Issues

This study was approved by Ethics Committee of University of Social Welfare and Rehabilitation Sciences (code of ethics: IR.USWR.REC.1396.275).

### Financial Support

This study was supported financially by Rofeideh Rehabilitation Hospital, University of Social Welfare and Rehabilitation Sciences, Tehran, Iran.

### References

- Morris SL, O'Sullivan PB, Murray KJ, Bear N, Hands B, Smith AJ. Hypermobility and Musculoskeletal Pain in Adolescents. *J Pediatr.* 2017;181:213-221. doi:10.1016/j.jpeds.2016.09.060
- Steinberg N, Hershkovitz I, Zeev A, Rothschild B, Sievner I. Joint Hypermobility and Joint Range of Motion in Young Dancers. *J Clin Rheumatol.* 2016;22(4):171-178. doi:10.1097/rhu.0000000000000420
- Ye S, Jing Q, Wei C, Lu J. Risk factors of non-specific neck pain and low back pain in computer-using office workers in China: a cross-sectional study. *BMJ Open.* 2017;7(4):e014914. doi:10.1136/bmjopen-2016-014914
- Karlsson L, Gerdle B, Takala EP, Andersson G, Larsson B. Associations between psychological factors and the effect of home-based physical exercise in women with chronic neck and shoulder pain. *SAGE Open Med.* 2016;4:2050312116668933. doi:10.1177/2050312116668933
- Pacey V, Nicholson LL, Adams RD, Munn J, Munns CF. Generalized joint hypermobility and risk of lower limb joint injury during sport: a systematic review with meta-analysis. *Am J Sports Med.* 2010;38(7):1487-1497. doi:10.1177/0363546510364838
- Remvig L, Jensen DV, Ward RC. Epidemiology of general joint hypermobility and basis for the proposed criteria for benign joint hypermobility syndrome: review of the literature. *J Rheumatol.* 2007;34(4):804-809.
- Saremi H, Yavarikia A, Jafari N. Generalized Ligamentous Laxity: An Important Predisposing Factor for Shoulder Injuries in Athletes. *Iran Red Crescent Med J.* 2016;18(6):e38903. doi:10.5812/ircmj.38903
- Russek LN, Errico DM. Prevalence, injury rate and, symptom frequency in generalized joint laxity and joint hypermobility syndrome in a "healthy" college population. *Clin Rheumatol.* 2016;35(4):1029-1039. doi:10.1007/s10067-015-2951-9
- Sohrbeck-Nohr O, Kristensen JH, Boyle E, Remvig L, Juul-Kristensen B. Generalized joint hypermobility in childhood is a possible risk for the development of joint pain in adolescence: a cohort study. *BMC Pediatr.* 2014;14:302. doi:10.1186/s12887-014-0302-7
- Hakim A, Grahame R. Joint hypermobility. *Best Pract Res Clin Rheumatol.* 2003;17(6):989-1004. doi:10.1016/j.berh.2003.08.001
- Trudelle-Jackson E, Leonard D, Morrow JR Jr. Musculoskeletal risk factors as predictors of injury in community-dwelling women. *Med Sci Sports Exerc.* 2014;46(9):1752-1757. doi:10.1249/mss.0000000000000295
- Bin Abd Razak HR, Bin Ali N, Howe TS. Generalized ligamentous laxity may be a predisposing factor for musculoskeletal injuries. *J Sci Med Sport.* 2014;17(5):474-478. doi:10.1016/j.jsams.2013.11.001
- Lugo R, Kung P, Ma CB. Shoulder biomechanics. *Eur J Radiol.* 2008;68(1):16-24. doi:10.1016/j.ejrad.2008.02.051
- Kibler WB, Sciascia A. Current concepts: scapular dyskinesis. *Br J Sports Med.* 2010;44(5):300-305. doi:10.1136/bjism.2009.058834
- Kibler WB, Sciascia A, Wilkes T. Scapular dyskinesis and its relation to shoulder injury. *J Am Acad Orthop Surg.* 2012;20(6):364-372. doi:10.5435/jaaos-20-06-364
- Arvidsson I, Gremark Simonsen J, Dahlqvist C, et al. Cross-sectional associations between occupational factors and musculoskeletal pain in women teachers, nurses and sonographers. *BMC Musculoskelet Disord.* 2016;17:35. doi:10.1186/s12891-016-0883-4
- Huang TS, Ou HL, Huang CY, Lin JJ. Specific kinematics and associated muscle activation in individuals with scapular dyskinesis. *J Shoulder Elbow Surg.* 2015;24(8):1227-1234. doi:10.1016/j.jse.2014.12.022
- Tate A, Turner GN, Knab SE, Jorgensen C, Strittmatter A, Michener LA. Risk factors associated with shoulder pain and disability across the lifespan of competitive swimmers. *J Athl Train.* 2012;47(2):149-158. doi:10.4085/1062-6050-47.2.149
- Warner JJ, Micheli LJ, Arslanian LE, Kennedy J, Kennedy R. Scapulothoracic motion in normal shoulders and shoulders with glenohumeral instability and impingement syndrome. A study using Moire topographic analysis. *Clin Orthop Relat Res.* 1992(285):191-199.
- Scheper M, de Vries J, Beelen A, de Vos R, Nollet F, Engelbert R. Generalized joint hypermobility, muscle strength and physical function in healthy adolescents and young adults. *Curr Rheumatol Rev.* 2014;10(2):117-125.
- Smith TO, Jerman E, Easton V, et al. Do people with benign joint hypermobility syndrome (BJHS) have reduced joint proprioception? A systematic review and meta-analysis.

- Rheumatol Int. 2013;33(11):2709-2716. doi:10.1007/s00296-013-2790-4
22. Sahin N, Baskent A, Ugurlu H, Berker E. Isokinetic evaluation of knee extensor/flexor muscle strength in patients with hypermobility syndrome. *Rheumatol Int.* 2008;28(7):643-648. doi:10.1007/s00296-007-0493-4
  23. Grahame R, Bird HA, Child A. The revised (Brighton 1998) criteria for the diagnosis of benign joint hypermobility syndrome (BJHS). *J Rheumatol.* 2000;27(7):1777-1779.
  24. Uhl TL, Kibler WB, Gecewich B, Tripp BL. Evaluation of clinical assessment methods for scapular dyskinesis. *Arthroscopy.* 2009;25(11):1240-1248. doi:10.1016/j.arthro.2009.06.007
  25. McClure P, Tate AR, Kareha S, Irwin D, Zlupko E. A clinical method for identifying scapular dyskinesis, part 1: reliability. *J Athl Train.* 2009;44(2):160-164. doi:10.4085/1062-6050-44.2.160
  26. Ghai S, Driller M, Ghai I. Effects of joint stabilizers on proprioception and stability: A systematic review and meta-analysis. *Phys Ther Sport.* 2017;25:65-75. doi:10.1016/j.ptsp.2016.05.006
  27. Chahal J, Leiter J, McKee MD, Whelan DB. Generalized ligamentous laxity as a predisposing factor for primary traumatic anterior shoulder dislocation. *J Shoulder Elbow Surg.* 2010;19(8):1238-1242. doi:10.1016/j.jse.2010.02.005
  28. Scheper MC, de Vries JE, Juul-Kristensen B, Nollet F, Engelbert RH. The functional consequences of generalized joint hypermobility: a cross-sectional study. *BMC Musculoskelet Disord.* 2014;15:243. doi:10.1186/1471-2474-15-243
  29. Jindal P, Narayan A, Ganesan S, MacDermid JC. Muscle strength differences in healthy young adults with and without generalized joint hypermobility: a cross-sectional study. *BMC Sports Sci Med Rehabil.* 2016;8:12. doi:10.1186/s13102-016-0037-x
  30. Jeremiah HM, Alexander CM. Do hypermobile subjects without pain have alteration to the feedback mechanisms controlling the shoulder girdle? *Musculoskeletal Care.* 2010;8(3):157-163. doi:10.1002/msc.178
  31. Fatoye F, Palmer S, Macmillan F, Rowe P, van der Linden M. Proprioception and muscle torque deficits in children with hypermobility syndrome. *Rheumatology (Oxford).* 2009;48(2):152-157. doi:10.1093/rheumatology/ken435
  32. Paletta GA Jr, Warner JJ, Warren RF, Deutsch A, Altchek DW. Shoulder kinematics with two-plane x-ray evaluation in patients with anterior instability or rotator cuff tearing. *J Shoulder Elbow Surg.* 1997;6(6):516-527. doi:10.1016/S1058-2746(97)90084-7
  33. Ogston JB, Ludewig PM. Differences in 3-dimensional shoulder kinematics between persons with multidirectional instability and asymptomatic controls. *Am J Sports Med.* 2007;35(8):1361-1370. doi:10.1177/0363546507300820
  34. Cameron KL, Duffey ML, DeBerardino TM, Stoneman PD, Jones CJ, Owens BD. Association of generalized joint hypermobility with a history of glenohumeral joint instability. *J Athl Train.* 2010;45(3):253-258. doi:10.4085/1062-6050-45.3.253
  35. Nodehi Moghadam A, Salimee MM. A comparative study on scapular static position between females with and without generalized joint hyper mobility. *Med J Islam Repub Iran.* 2012;26(3):97-102.
  36. Alibazi RJ, Moghadam AN, Cools AM, Bakhshi E, Ahari AA. The Effect of Shoulder Muscle Fatigue on Acromiohumeral Distance and Scapular Dyskinesis in Women With Generalized Joint Hypermobility. *J Appl Biomech.* 2017;33(6):424-430. doi:10.1123/jab.2016-0056
  37. Ebaugh DD, McClure PW, Karduna AR. Effects of shoulder muscle fatigue caused by repetitive overhead activities on scapulothoracic and glenohumeral kinematics. *J Electromyogr Kinesiol.* 2006;16(3):224-235. doi:10.1016/j.jelekin.2005.06.015
  38. Tsai NT, McClure PW, Karduna AR. Effects of muscle fatigue on 3-dimensional scapular kinematics. *Arch Phys Med Rehabil.* 2003;84(7):1000-1005. doi:10.1016/S0003-9993(03)00127-8
  39. Chopp JN, Fischer SL, Dickerson CR. The specificity of fatiguing protocols affects scapular orientation: Implications for subacromial impingement. *Clin Biomech (Bristol, Avon).* 2011;26(1):40-45. doi:10.1016/j.clinbiomech.2010.09.001

**Copyright** © 2018 The Author(s); This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.